## Mold and Mycobacteria Blood Collection Instructions



## This sample is a blood culture and should be drawn in the same way as blood cultures! Peripheral stick only and do NOT draw from line!

Kit contents: BACTEC MycoF/Lytic Bottle, ChloraPrep solultion (CHG), Alcohol Wipe

Test Name	Test Code	Collection Device	Preferred Volume	Minimum Volume
Mold Blood Culture	LAB242	BACTEC MycoF/Lytic	5 mL	1 mL
Mycobacteria Blood Culture	LAB246	BACTEC MycoF/Lytic	5 mL	1 mL

## **Collection Procedure:**

- 1. Prep peripheral stick site with ChloraPrep solution (CHG) using **back-and-forth** friction.
- 2. Scrub for 30 seconds. Allow the CHG on the skin to dry.
- 3. While skin is drying, disinfect rubber top with alcohol wipe.
- 4. Allow the disinfectant to dry completely.
- 5. Aseptically draw blood 5mL into the bottle using the 5 mL graduation marks on the bottle label (see image, left).
- 6. Ensure BACTEC MycoF/Lytic Bottle is labeled appropriately (see image, right).
- 7. Maintain sterility and forward promptly to the laboratory.

## <u>NOTE:</u> One bottle required per test ordered. If both Mold blood and Mycobacteria blood cultures are ordered, 2 BACTEC MycoF/Lytic bottles are required.

Call BJH Microbiology lab with questions: 314-362-3898.



How to Properly Label:

